



Fisheries and Oceans
Canada

Pêches et Océans
Canada

Ecosystems and
Oceans Science

Sciences des écosystèmes
et des océans

Canadian Science Advisory Secretariat (CSAS)

Research Document 2016/058

Québec Region

Physical Oceanographic Conditions in the Gulf of St. Lawrence in 2015

P. S. Galbraith¹, J. Chasse², C. Caverhill³, P. Nicot⁴, D. Gilbert¹,
B. Pettigrew¹, D. Lefebvre¹, D. Brickman¹, L. Devine¹, C. Lefleur¹

¹ Fisheries and Oceans Canada, Québec Region,
Maurice Lamontagne Institute,
P.O. Box 10300, Mont-Joli, Québec, G5H 3Z4

² Fisheries and Oceans Canada, Gulf Region,
P.O. Box 5020, Moncton, New Brunswick, E1C 9B6

³ Fisheries and Oceans Canada, Maritimes Region,
Bedford Institute of Oceanography
P.O. Box 1010, Dartmouth, Nova Scotia, B2Y 4A2

⁴ Institut des sciences de la mer de Rimouski
Université du Québec à Rimouski
210 allée des Ursulines, Rimouski, Québec, G5L 3A1

Foreword

This series documents the scientific basis for the evaluation of aquatic resources and ecosystems in Canada. As such, it addresses the issues of the day in the time frames required and the documents it contains are not intended as definitive statements on the subjects addressed but rather as progress reports on ongoing investigations.

Research documents are produced in the official language in which they are provided to the Secretariat.

Published by:

Fisheries and Oceans Canada
Canadian Science Advisory Secretariat
200 Kent Street
Ottawa ON K1A 0E6

<http://www.dfo-mpo.gc.ca/csas-sccs/>
csas-sccs@dfo-mpo.gc.ca



© Her Majesty the Queen in Right of Canada, 2016
ISSN 1919-5044

Correct citation for this publication:

Galbraith, P.S., Chassé, J., Caverhill, C., Nicot, P., Gilbert, D., Pettigrew, B., Lefavre, D.,
Brickman, D., Devine, L., and Lafleur, C. 2016. Physical Oceanographic Conditions in the
Gulf of St. Lawrence in 2015. DFO Can. Sci. Advis. Sec. Res. Doc. 2016/056. v + 90 p.

TABLE OF CONTENTS

ABSTRACT.....	IV
RÉSUMÉ.....	V
INTRODUCTION.....	1
AIR TEMPERATURE.....	2
PRECIPITATION AND FRESHWATER RUNOFF.....	2
SURFACE LAYER.....	3
SST SEASONAL CYCLE CLIMATOLOGY.....	4
SST IN 2015.....	4
SEA ICE.....	6
WINTER WATER MASSES.....	8
COLD INTERMEDIATE LAYER.....	9
PREDICTION FROM THE MARCH SURVEY.....	9
AUGUST CIL BASED ON THE MULTI-SPECIES SURVEY.....	9
NOVEMBER CIL CONDITIONS IN THE ST. LAWRENCE ESTUARY.....	10
GILBERT AND PETTIGREW (1997) CIL INDEX.....	10
MAGDALEN SHALLOWS JUNE SURVEY.....	11
BOTTOM WATER TEMPERATURES ON THE MAGDALEN SHALLOWS.....	11
DEEP WATERS (>150 M).....	12
BOTTOM WATER TEMPERATURES IN AUGUST AND SEPTEMBER.....	12
TEMPERATURE AND SALINITY MONTHLY MEANS.....	13
DISSOLVED OXYGEN AND HYPOXIA IN THE ST. LAWRENCE ESTUARY.....	13
SEASONAL AND REGIONAL AVERAGE TEMPERATURE STRUCTURE.....	14
CURRENTS AND TRANSPORTS.....	15
HIGH FREQUENCY SAMPLING AZMP STATIONS.....	16
OUTLOOK FOR 2015.....	16
SUMMARY.....	17
KEY FINDINGS.....	17
ACKNOWLEDGEMENTS.....	18
REFERENCES.....	19

ABSTRACT

An overview of physical oceanographic conditions in the Gulf of St. Lawrence (GSL) in 2015 is presented as part of the Atlantic Zone Monitoring Program (AZMP). AZMP data as well as data from regional monitoring programs are analysed and presented in relation to long-term means. The annual average freshwater runoff of the St. Lawrence River measured at Québec City was below normal and its combination with rivers flowing into the Estuary (RIVSUM II) was about normal in 2015 (-1.0 SD and +0.1 SD respectively). The spring freshet was below-normal at -1.1 SD and -0.5 SD, but its timing was normal. The cold winter of 2015 created a thick surface mixed layer over the GSL with near-freezing temperatures, but a sea ice cover that was only slightly above-normal. The August cold intermediate layer (CIL) showed warmer minimum temperature (+0.8 SD) and thinner volume colder than 1°C (-1.5 SD) than normal in spite of the cold winter conditions. Including the colder temperatures from the June survey led to a near-normal CIL minimum temperature index (+0.3 SD). Sea-surface temperatures averaged over the Gulf were generally below normal until July and above normal from August to September 2015, leading to a near-normal May-November average (+0.3°C, +0.5 SD). Record highs were nevertheless reached in September averaged over the Gulf (+1.8°C, +2.3 SD) and in the following regions: Estuary (+1.9°C, +2.3 SD), Northwest Gulf (+1.9°C, +1.9 SD), Anticosti Channel (+2.1°C, +2.1 SD) and Magdalen Shallows (+1.4°C, +2.0 SD). The timing of summer warming onset was near-normal while fall cooling was later than normal (+1.7 weeks, +1.4 SD). Deep water temperatures have been increasing overall in the Gulf, with inward advection from Cabot Strait where temperature had reached a record high (since 1915) in 2012 at 200 m. Temperature averaged over the Gulf at depths from 150 to 300 m all show record highs in 2015 and attained values above 6°C at 250 and 300 m for the first time since 1915. The bottom area covered by waters warmer than 6°C increased in 2015 in Anticosti Channel, Esquiman Channel and Central Gulf, reaching a record high value in Anticosti Channel and Esquiman Channel while reducing its bottom habitat area in the temperature range of 5–6°C.

Conditions océanographiques physiques dans le golfe du Saint-Laurent en 2015

RÉSUMÉ

Le présent document donne un aperçu des conditions d'océanographie physique qui ont prévalu dans le golfe du Saint-Laurent en 2015 et est un produit du Programme de monitoring de la zone Atlantique (PMZA). Les données du PMZA ainsi que de programmes de monitoring régionaux sont analysées et présentées en relation avec des moyennes à long terme. Les débits du fleuve Saint-Laurent et de l'indice RIVSUM II étaient respectivement sous et près de la normale en 2015 (- 1,0 É.T. et + 0,1 É.T. respectivement). La crue printanière était sous la normale (- 1,1 É.T. et - 0,5 É.T.) mais sa phénologie était normale. L'hiver froid de 2015 a causé la formation d'une volumineuse couche de surface avec des températures près du point de congélation, par contre le couvert de glace de mer n'a atteint un volume saisonnier maximal que légèrement au-dessus de la normale. La couche intermédiaire froide (CIF) du mois d'août était chaude (+ 0,8 É.T.) et mince (- 1,5 É.T. pour le volume des eaux sous 1 °C) en dépit de l'hiver froid. L'inclusion des données plus froides de juin mène à l'évaluation d'une CIF près de la normale (+ 0,3 É.T.). Les températures de l'eau à la surface de étaient généralement sous la normale jusqu'en juillet, et au-dessus de la normale en août et septembre, conduisant à une moyenne de mai à novembre qui a été près de la normale (+ 0,3 °C, + 0,5 É.T.). Le mois de septembre a néanmoins connu un record de + 1,8 °C au-dessus de la climatologie (+ 2,3 É.T.). Des records ont aussi été battus dans les régions suivantes : estuaire (+ 1,9 °C, + 2,3 É.T.), nord-ouest du golfe (+ 1,9 °C, + 1,9 É.T.), chenal Anticosti (+ 2,1 °C, + 2,1 É.T.) et le Plateau madelinien (+ 1,4 °C, + 2,0 É.T.). Le réchauffement printanier est survenu à une date normale, tandis que le refroidissement d'automne a été tardif (+ 1,7 semaine, + 1,4 É.T.). Les températures des eaux profondes du golfe ont depuis quelques années été en augmentation avec le transport depuis le détroit de Cabot d'eaux qui avaient atteint une température record (depuis 1915) en 2012 à 200 m. Globalement, les températures de 150 m à 300 m de profondeur ont atteint un record de série et dépassent le seuil de 6 °C pour la première fois à 250 et 300 m depuis 1915. La superficie du fond marin recouvert par des températures plus grandes que 6 °C a augmenté dans le chenal d'Anticosti, le chenal Esquiman ainsi que dans le centre du golfe, atteignant un record de série dans le chenal d'Anticosti ainsi que dans le chenal Esquiman au détriment de l'habitat de fond dans la plage de température de 5 à 6 °C.

INTRODUCTION

This document examines the physical oceanographic conditions and related atmospheric forcing in the Gulf of St. Lawrence in 2015 (Figure 1). It complements similar reviews of the environmental conditions on the Newfoundland and Labrador Shelf and the Scotian Shelf and Gulf of Maine as part of the Atlantic Zone Monitoring Program (AZMP; see Theriault et al., 1998 for background information on the program and Colbourne et al. 2015, and Hebert et al. 2015 for examples of past reviews in other AZMP regions). The last detailed report of physical oceanographic conditions in the Gulf of St. Lawrence was produced for the year 2014 (Galbraith et al. 2015). Specifically, it discusses air temperature, freshwater runoff, sea-ice volume, surface water temperature and salinity, winter water mass conditions (e.g., the near-freezing mixed layer volume, the volume of dense water that entered the Gulf through the Strait of Belle Isle), the summertime cold intermediate layer (CIL), and the temperature, salinity, and dissolved oxygen of the deeper layers. Some of the variables are spatially averaged over distinct regions of the Gulf (Figure 2). The report uses data obtained from the Department of Fisheries and Oceans' (DFO) Atlantic Zone Monitoring Program (AZMP), other DFO surveys, and other sources. Environmental variables are usually expressed as anomalies, i.e., deviations from their long-term mean. The long-term mean or normal conditions are calculated for the standard 1981–2010 reference period when possible. Furthermore, because these series have different units ($^{\circ}\text{C}$, m^3 , m^2 , etc.), each anomaly time series is normalized by dividing by its standard deviation (SD), also calculated for the standard reference when possible. This allows a more direct comparison of the various series. Missing data are represented by grey cells in the tables, values within ± 0.5 SD of the average as white cells, and conditions corresponding to warmer than normal (higher temperatures, reduced ice volumes, reduced cold-water volumes or areas) by more than 0.5 SD as red cells, with more intense reds corresponding to increasingly warmer conditions. Similarly, blue represents colder than normal conditions. Higher than normal freshwater inflow is shown as red, but does not necessarily correspond to warmer-than-normal conditions. Higher than normal stratification values are shown in blue because they are usually caused by lower upper layer salinity.

The summertime water column in the Gulf of St. Lawrence consists of three distinct layers: the surface layer, the cold intermediate layer (CIL), and the deeper water layer (Figure 3). Surface temperatures typically reach maximum values in early to mid-August. Gradual cooling occurs thereafter, and wind forced mixing during the fall leads to a progressively deeper and cooler mixed layer, eventually encompassing the CIL. During winter, the surface layer thickens partly because of buoyancy loss (cooling and reduced runoff) and brine rejection associated with sea-ice formation, but mostly from wind-driven mixing prior to ice formation (Galbraith 2006). The surface winter layer extends to an average depth of 75 m but >150 m in places such as the Mécatina Trough (where intruding waters through the Strait of Belle Isle from the Labrador Shelf may extend from the surface to the bottom in depths >200 m) by the end of March when temperatures have decreased to near freezing (-1.8 to 0°C) (Galbraith 2006). During spring, surface warming, sea-ice melt waters, and continental runoff produce a lower-salinity and higher-temperature surface layer. Underneath this surface layer, cold waters from the previous winter are partly isolated from the atmosphere and form the summer CIL. This layer will persist until the next winter, gradually warming up and deepening during summer (Gilbert and Pettigrew 1997; Cyr et al. 2011) and more rapidly during the fall as vertical mixing intensifies.

This report considers these three layers in turn but first air temperature is examined because it is a significant driver of the surface layer, followed by the freshwater runoff. The winter sea ice and winter oceanographic conditions are described; these force the summer CIL, which is

presented next. The deeper waters, mostly isolated from exchanges with the surface, are presented last along with a summary of major oceanographic surveys.

AIR TEMPERATURE

The air temperature data are the second generation of homogenized surface air temperature data, part of the Adjusted and Homogenized Canadian Climate Data (AHCCD), which accounts for shifts due to the relocation of stations, changes in observing practices and automation (Vincent et al. 2012). The monthly air temperature anomalies for several stations around the Gulf are shown in Figure 4 for 2014 and 2015, as well as the average of all station anomalies.

Figure 5 shows the annual, winter (December-March), and April-November mean air temperature anomalies averaged over all available stations shown in Figure 4 since 1873. Record-high annual and winter temperatures occurred in 2010 and record-high April-November temperatures in 2012. Galbraith et al. (2012) found the average April-November air temperature over the Gulf from Environment Canada's National Climate Data and Information Archive (NCDIA) to be a good proxy for May-November sea-surface temperature over the Gulf (but excluding the estuary) and found within the former a warming trend of 0.9°C per century between 1873 and 2011; the same trend is found here over the selected AHCCD stations between 1873 and 2015 (Figure 5). The NCDIA December-March air temperatures in the western Gulf were found to be highly correlated ($r^2=0.67$) with sea-ice properties, as well as with winter mixed layer volumes (Galbraith et al. 2010). Galbraith et al. (2013) found slightly higher correlations ($r^2=0.72$) with sea-ice using December-February AHCCD averages, possibly because March temperature are of less importance during low sea-ice cover since much of the sea-ice cover decrease has occurred much earlier in February.

Air temperatures were well below normal in February 2015, by 6.5°C at Mont-Joli (coldest since 1923, -2.4 SD), by 7.3°C at Bathurst (1922, -2.7 SD), by 5.1°C at Charlottetown (1923, -2.4 SD), by 5.4°C at Gaspé (1939, -2.0 SD) and by 4.4°C on average over available Gulf stations (coldest since 1993 and 15th lowest since 1873 at -1.6 SD). Overall the winter was cold, with a December-March average air temperature anomaly of -1.5°C (-1.0 SD). August air temperatures were at series record highs at Natashquan (since 1915, +2.9°C, +2.9 SD), Baie-Comeau (since 1965, +2.5°C, +2.6 SD), Mont-Joli (since 1877, +2.3°C, +2.0 SD) and Charlottetown (since 1873, +2.8°C, +3.0 SD). Averaged over all stations, August had the second warmest (after 2012) air temperatures on record (since 1873, +2.2°C, +2.8 SD). April-November air temperatures were normal (anomaly of 0.0°C) and the annual averaged air temperature anomaly was near-normal at -0.4°C overall (-0.4 SD).

PRECIPITATION AND FRESHWATER RUNOFF

Freshwater runoff data for the St. Lawrence River are updated monthly (Figure 6, lower curve) using the water level method from Bourgault and Koutitonsky (1999) and are available from the St. Lawrence Global Observatory. A hydrological watershed model was used to estimate the monthly runoff since 1948 for all other major rivers flowing into the Gulf of St. Lawrence, with discharge locations as shown in Figure 7. The precipitation data (NCEP reanalysis, six hourly intervals) used as input in the model were obtained from the NOAA-CIRES Climate Diagnostics Center (Boulder, Colorado, USA; Kalnay et al. 1996). The data were interpolated to a 1/4° resolution grid and the water routed to river mouths using a simple algorithm described here. When air temperatures were below freezing, the water was accumulated as snow in the watershed and later melted as a function of warming temperatures. Water regulation is modelled for three rivers that flow into the estuary (Saguenay, Manicouagan, Outardes) for which the annual runoff is redistributed following the climatology of the true regulated runoffs for

12 months thereafter. Runoffs were summed for each region shown and the climatology established for the 1981–2010 period. The waters that flow into the Estuary (region 1, Figure 7) were added to the St. Lawrence River runoff measured at Quebec City to produce the RivSUM II index, although no advection lags were introduced (Figure 6, upper curve).

Monthly anomalies of the summed runoffs for 2014 and 2015 are shown in Figure 6. Rivers other than the St. Lawrence contribute about $5\,000\text{ m}^3\text{ s}^{-1}$ runoff to the Estuary, the equivalent of 40% of the St. Lawrence River, while the other tributaries distributed along the border of the GSL provide an additional $3\,500\text{ m}^3\text{ s}^{-1}$ in freshwater runoff to the system. River regulation has a strong impact on the relative contributions of sources. For example, in May 2015 the higher-than-average river runoff into the Estuary (an effect of the heavy precipitation in 2014 and river regulation) was almost as important as the below-normal St. Lawrence run-off. The 2015 simulation shows that rivers in regions 2, 3, 4 and 5 behaved similarly, with below normal runoff in winter and above normal in June, 2015, through to August in most regions. The long-term time series are shown, summed by large basins, in Figure 9. Broad long-term patterns of runoff over the large basins were similar to that of the St. Lawrence River but interannual variability is low in the Northeast basin and Magdalen Shallows basin. The average run-off into the Estuary was above-normal at +2.7 SD, the highest since 1964. The annual average runoff of the St. Lawrence River measured at Quebec City and RivSUM II both show a general downward trend from the mid-1970s until 2001, an upwards trend between 2001 and 2011 and were respectively below-normal and normal in 2015 (Figure 9) at $11\,100\text{ m}^3\text{ s}^{-1}$ (-1.0 SD) and $17\,000\text{ m}^3\text{ s}^{-1}$ (+0.6 SD). Runoff was low from March through May, consistent with the very cold air temperatures. The spring freshet was below-normal at -1.1 SD in April for the St. Lawrence River and -0.5 SD in May for RivSUM II, but its timing was normal (Figure 6). The fall freshet was also above normal and later than normal.

SURFACE LAYER

The surface layer conditions of the Gulf are monitored by several complementary methods. The shipboard thermosalinograph network (Galbraith et al. 2002) consists of temperature-salinity sensors (SBE-21, Sea-Bird Electronics Inc., Bellevue, WA) that have been installed on various ships starting with the commercial ship Cicero of Oceanex Inc. in 1999 (retired in 2006) and on the Cabot from 2006 to fall 2013. The Oceanex Cormaigra, was outfitted with a thermosalinograph in early 2015.

The second data source is the thermograph network (Gibert et al. 2004, Galbraith et al. 2007), which consists of a number of stations with moored instruments recording water temperature every 30 minutes (Figure 10). Most instruments are installed on Coast Guard buoys that are deployed in the ice-free season, but a few stations are monitored year-round. The data are typically only available after the instruments are recovered except for oceanographic buoys that transmit data in real-time. Data from Shediac station acquired by the DFO Gulf Region are also included here. The network provides valuable near-surface temperature data at fixed sites and at short sampling intervals, but usually not in real-time nor during winter months.

The third data source are 1 km resolution monthly composite Sea Surface Temperature (SST) generated using National Oceanic and Atmospheric Administration (NOAA) and European Organisation for the Exploitation of Meteorological Satellites (EUMETSAT) Advanced Very High Resolution Radiometer (AVHRR) satellite images available from the Maurice Lamontagne Institute sea surface temperature processing facility (details in Galbraith and Larueche 2011, and Galbraith et al. 2012). These data are available for the period of 1985–2013. AVHRR composites of 1.5-km resolution provided by the Bedford Institute of Oceanography (BIO) Operational Remote Sensing group complete the data set. Monthly climatologies for the period

of 1985–2012 version to both products were compared at the 1.5 km grid level for cross calibration. The SIO product is adjusted to the ISU product climatology as $SST_{adj} = SST_{SIO} + 0.13 + 0.13^{\circ}\text{C}$ adjustment at 2°C , -0.54°C at 20°C). This is in part explained by the fact that the ISU product used all available SST images while the SIO product uses only daytime passes, introducing a slight diurnal bias. This adjustment has decreased somewhat the 2010 record high temperatures reported in Goffarth et al. 2015.

SST SEASONAL CYCLE CLIMATOLOGY

The May to November cycle of weekly averaged surface temperature is illustrated in Figure 11 using a 1985–2010 climatology based on AVHRR remote sensing data for ice-free months complemented by 2007–2010 thermistograph data for the winter months. Goffarth et al. (2012) have shown that Gulf averaged air temperature and SST monthly climatologies match up quite well with SST lagging air temperature by half a month. Maximum sea surface temperatures are reached on average during the second week of August but that can vary by a few weeks from year to year. The maximum surface temperature averages 15.0°C over the Gulf during the second week of August (1985–2010), but there are spatial differences: temperatures on the Magdalen Shallows are the warmest in the Gulf, averaging 16.1°C over that area, and the coldest are at the head of the St. Lawrence Estuary and upwelling areas along the coast north shore.

Figure 12 shows a mean annual cycle of water temperature at a depth of 5 m along the shoreline to St. John's shipping route based on thermistograph data collected from 2002 to 2015. Data were used from all instrumented ships that were within the main shipping route and to fill data gaps. The data were averaged for each day of the year at intervals of 0.1 degree of longitude to create a composite along the ship track. The most striking feature is the area at the head of the Laurentian Trough (55.5°N), where strong vertical mixing leads to cold summer water temperatures (around 5°C to 6°C and sometimes lower) and winter temperatures that are always above freezing (see also Figure 11). The climatology shows the progression to winter conditions, first reaching near-freezing temperatures in the Estuary and then progressing coastward with time, usually reaching Cabot Strait by the end of the winter.

Temperature anomaly time series and the 2005–2015 climatologies were constructed for selected sections that are crossed by the ship (Figure 13). Although the profiles are quite similar between the two sections, the near surface temperature climatology at Tadoussac (head of the Laurentian Trough) contrasts with that nearby in the Estuary, as noted above. Winter temperatures are on average 3.7°C warmer at the Tadoussac section; the maximum monthly mean temperature in summer is only 7.1°C compared with 8.8°C at the nearby Estuary section and up to 13.2°C at the Mont-Louis section. The table in Figure 13 provides a quick reference to the interannual near surface temperature variations at the selected sections as well as monthly averages for the year in route.

SST IN 2015

Thermistograph data show that near-freezing surface water conditions disappeared earlier than normal in the Estuary and lasted longer in the Gulf near Cabot Strait (Figure 12). The 2015 monthly mean sea surface temperatures from AVHRR imagery are shown in Figure 14 as colour-coded maps and the corresponding temperature anomaly maps are shown in Figure 15 (referenced to 1985–2010 monthly climatologies). Missing anomaly pixels are due to incomplete climatologies during the ice-covered winter months. The SST information is summarized in Figure 16, showing the 2015 monthly average temperatures versus the climatologies spatially averaged over the Gulf and over each of the eight regions delimited by the areas shown in Figure 2. The information also appears again in tabular form in Figure 17 with the same

the average height of the forest was 10.0 m. The average height of the forest was 10.0 m. The average height of the forest was 10.0 m.

The average height of the forest was 10.0 m. The average height of the forest was 10.0 m. The average height of the forest was 10.0 m.

The average height of the forest was 10.0 m. The average height of the forest was 10.0 m. The average height of the forest was 10.0 m.

The average height of the forest was 10.0 m. The average height of the forest was 10.0 m. The average height of the forest was 10.0 m.

The average height of the forest was 10.0 m. The average height of the forest was 10.0 m. The average height of the forest was 10.0 m.

The average height of the forest was 10.0 m. The average height of the forest was 10.0 m. The average height of the forest was 10.0 m.

Copyright © 2006 John Wiley & Sons, Ltd.

[illegible]

Figure 10 shows the log-likelihood surface for the parameters of the lognormal distribution. The surface is relatively flat, indicating that the lognormal distribution is a good fit for the data.

[illegible][illegible]

[illegible]

1. **Introduction**
 2. **Background**
 3. **Methodology**
 4. **Results**
 5. **Discussion**
 6. **Conclusion**
 7. **References**
 8. **Appendix**
 9. **Index**
 10. **Table of Contents**
 11. **Figure 1**
 12. **Figure 2**
 13. **Figure 3**
 14. **Figure 4**
 15. **Figure 5**
 16. **Figure 6**
 17. **Figure 7**
 18. **Figure 8**
 19. **Figure 9**
 20. **Figure 10**
 21. **Figure 11**
 22. **Figure 12**
 23. **Figure 13**
 24. **Figure 14**
 25. **Figure 15**
 26. **Figure 16**
 27. **Figure 17**
 28. **Figure 18**
 29. **Figure 19**
 30. **Figure 20**
 31. **Figure 21**
 32. **Figure 22**
 33. **Figure 23**
 34. **Figure 24**
 35. **Figure 25**
 36. **Figure 26**
 37. **Figure 27**
 38. **Figure 28**
 39. **Figure 29**
 40. **Figure 30**
 41. **Figure 31**
 42. **Figure 32**
 43. **Figure 33**
 44. **Figure 34**
 45. **Figure 35**
 46. **Figure 36**
 47. **Figure 37**
 48. **Figure 38**
 49. **Figure 39**
 50. **Figure 40**
 51. **Figure 41**
 52. **Figure 42**
 53. **Figure 43**
 54. **Figure 44**
 55. **Figure 45**
 56. **Figure 46**
 57. **Figure 47**
 58. **Figure 48**
 59. **Figure 49**
 60. **Figure 50**
 61. **Figure 51**
 62. **Figure 52**
 63. **Figure 53**
 64. **Figure 54**
 65. **Figure 55**
 66. **Figure 56**
 67. **Figure 57**
 68. **Figure 58**
 69. **Figure 59**
 70. **Figure 60**
 71. **Figure 61**
 72. **Figure 62**
 73. **Figure 63**
 74. **Figure 64**
 75. **Figure 65**
 76. **Figure 66**
 77. **Figure 67**
 78. **Figure 68**
 79. **Figure 69**
 80. **Figure 70**
 81. **Figure 71**
 82. **Figure 72**
 83. **Figure 73**
 84. **Figure 74**
 85. **Figure 75**
 86. **Figure 76**
 87. **Figure 77**
 88. **Figure 78**
 89. **Figure 79**
 90. **Figure 80**
 91. **Figure 81**
 92. **Figure 82**
 93. **Figure 83**
 94. **Figure 84**
 95. **Figure 85**
 96. **Figure 86**
 97. **Figure 87**
 98. **Figure 88**
 99. **Figure 89**
 100. **Figure 90**
 101. **Figure 91**
 102. **Figure 92**
 103. **Figure 93**
 104. **Figure 94**
 105. **Figure 95**
 106. **Figure 96**
 107. **Figure 97**
 108. **Figure 98**
 109. **Figure 99**
 110. **Figure 100**
 111. **Figure 101**
 112. **Figure 102**
 113. **Figure 103**
 114. **Figure 104**
 115. **Figure 105**
 116. **Figure 106**
 117. **Figure 107**
 118. **Figure 108**
 119. **Figure 109**
 120. **Figure 110**
 121. **Figure 111**
 122. **Figure 112**
 123. **Figure 113**
 124. **Figure 114**
 125. **Figure 115**
 126. **Figure 116**
 127. **Figure 117**
 128. **Figure 118**
 129. **Figure 119**
 130. **Figure 120**
 131. **Figure 121**
 132. **Figure 122**
 133. **Figure 123**
 134. **Figure 124**
 135. **Figure 125**
 136. **Figure 126**
 137. **Figure 127**
 138. **Figure 128**
 139. **Figure 129**
 140. **Figure 130**
 141. **Figure 131**
 142. **Figure 132**
 143. **Figure 133**
 144. **Figure 134**
 145. **Figure 135**
 146. **Figure 136**
 147. **Figure 137**
 148. **Figure 138**
 149. **Figure 139**
 150. **Figure 140**
 151. **Figure 141**
 152. **Figure 142**
 153. **Figure 143**
 154. **Figure 144**
 155. **Figure 145**
 156. **Figure 146**
 157. **Figure 147**
 158. **Figure 148**
 159. **Figure 149**
 160. **Figure 150**
 161. **Figure 151**
 162. **Figure 152**
 163. **Figure 153**
 164. **Figure 154**
 165. **Figure 155**
 166. **Figure 156**
 167. **Figure 157**
 168. **Figure 158**
 169. **Figure 159**
 170. **Figure 160**
 171. **Figure 161**
 172. **Figure 162**
 173. **Figure 163**
 174. **Figure 164**
 175. **Figure 165**
 176. **Figure 166**
 177. **Figure 167**
 178. **Figure 168**
 179. **Figure 169**
 180. **Figure 170**
 181. **Figure 171**
 182. **Figure 172**
 183. **Figure 173**
 184. **Figure 174**
 185. **Figure 175**
 186. **Figure 176**
 187. **Figure 177**
 188. **Figure 178**
 189. **Figure 179**
 190. **Figure 180**
 191. **Figure 181**
 192. **Figure 182**
 193. **Figure 183**
 194. **Figure 184**
 195. **Figure 185**
 196. **Figure 186**
 197. **Figure 187**
 198. **Figure 188**
 199. **Figure 189**
 200. **Figure 190**
 201. **Figure 191**
 202. **Figure 192**
 203. **Figure 193**
 204. **Figure 194**
 205. **Figure 195**
 206. **Figure 196**
 207. **Figure 197**
 208. **Figure 198**
 209. **Figure 199**
 210. **Figure 200**
 211. **Figure 201**
 212. **Figure 202**
 213. **Figure 203**
 214. **Figure 204**
 215. **Figure 205**
 216. **Figure 206**
 217. **Figure 207**
 218

1. The first step is to identify the problem. This involves understanding the situation and the goals that need to be achieved.

1. The first step in the process of creating a new product is to identify a market need. This involves conducting market research to understand the current market landscape, identify gaps, and determine the target audience.

1. **Introduction:** The purpose of this study is to investigate the impact of social media on the mental health of adolescents. The study aims to explore the relationship between social media usage and various mental health outcomes, including self-esteem, anxiety, and depression.

© 2000 Blackwell Science Ltd, *Journal of Internal Medicine* 247: 103–110

(Figure 23). Deep temperatures in the Strait of Belle Isle were below -1°C up to May 20th (Figure 20), which could be associated with prolonged entry of Labrador Shelf waters into the Gulf.

The cold mixed layer depth typically reaches about 75 m in the Gulf and is usually delimited by the -1°C isotherm because the mixed layer is typically near-freezing and deeper waters are much warmer (Galbraith 2006). In March 2010 and 2011 much of the mixed layer was warmer than -1°C such that the criterion of $T < 0^{\circ}\text{C}$ was also introduced (see middle panels of Figure 41). The cold surface layer is the product of local formation as well as cold waters advected from the Labrador Shelf, and can consist either of a single water mass or of layers of increasing salinity with depth. Integrating the cold layer depth over the area of the Gulf (excluding the Estuary and the Strait of Belle Isle) yields a cold-water ($< -1^{\circ}\text{C}$) volume of 13 500 km³ in 2015 (Figure 43), 0.5 SD above the 1996–2015 average and similar to the volumes observed in 2001, 2002 and 2004. The time series of winter cold-water ($< -1^{\circ}\text{C}$) volume observed in the Gulf is shown in Figure 43. The mixed layer volume increases only to 16 200 km³ when water temperatures $< 1^{\circ}\text{C}$ are considered which is close to the 1996–2015 average. This total volume of cold water corresponds to 62% of the total water volume of the Gulf (22 000 km³, excluding the Estuary).

COLD INTERMEDIATE LAYER

PREDICTION FROM THE MARCH SURVEY

The summer OI minimum temperature index (Gibert and Pettigrew 1997) has been found to be tightly correlated with the Gulf (excluding the estuary) volume of cold water ($< -1^{\circ}\text{C}$) measured the previous March when much of the mixed layer is near-freezing (Galbraith 2006, updated relation in right panel of Figure 43). This is expected because the OI is the remnant of the winter cold surface layer. A measurement of the volume of cold water present in March is therefore a valuable tool for forecasting the coming summer OI conditions. The winter mixed layer in 2015 was thick and near-freezing throughout the Gulf, up to Cabot Strait, after four consecutive winters with warmer conditions. The overall thickness and volume of the layer colder than -1°C was 0.5 SD above normal in 2015 in spite of the record low volume of Labrador Shelf water that usually also contributes towards the overall volume. The Cold Intermediate Layer for summer 2015 was therefore forecasted to be a bit warmer than in 2014, with a Gilbert and Pettigrew (1997) index of around -0.8°C compared to -0.5°C in 2014 (Galbraith et al. 2015).

AUGUST OI BASED ON THE MULTI-SPECIES SURVEY

The OI minimum temperature, thickness and volume for $T < 0^{\circ}\text{C}$ and $< 1^{\circ}\text{C}$ were estimated using temperature profiles from all sources for August and September. Most data are from the multi-species surveys in September for the Magdalen Shallows and August for the rest of the Gulf. Using all available temperature profiles, each 1-m depth layer of the Gulf was spatially interpolated for temperature, with the interpolated field bound between the minimum and maximum values observed within each of the different regions of the Gulf (Figure 2) to avoid spurious extrapolations. The OI thickness at each grid point is simply the sum of depth bins below the threshold temperature, and the OI minimum temperature is only defined at grid points where temperature rose by at least 0.5 $^{\circ}\text{C}$ at depths greater than that of the minimum, or if the grid point minimum temperature is below the OI spatial average of the Gulf.

Figure 44 shows the gridded interpolation of the OI thickness $< 1^{\circ}\text{C}$ and $< 0^{\circ}\text{C}$ and the OI minimum temperature for August–September 2015 as well their 1996–2015 climatology (1996–2015 for Magdalen Trough). The OI thickness for $T < 0^{\circ}\text{C}$ and $T < 1^{\circ}\text{C}$ decreased to below-normal values since 2014, with conditions similar to 2013 (not shown). Similar maps were produced for

all years back to 1971 (although some years have no data in some regions), allowing the calculation of volumes for each region for each year as well as the climatologies shown on the left side of Figure 44. The 2015 CIL water mass was thinner and warmer than the 1985-2010 climatologies, with particularly warm conditions in the Estuary, with minimum temperatures above 1°C in the western half.

The time series of the regional August-September CIL volumes are shown in Figure 45 (for <0°C and <1°C). All regions except Cabot Strait show decreased CIL volumes in 2015 compared to 2014. Figure 46 shows the total volume of CIL water (<0°C and <1°C) and the average CIL core temperature from the August-September interpolated grids (e.g., Figure 44). The CIL areal minimum temperature average and volume shown in Figure 46 exclude data from the Magdalen Trough which has very different water masses from the rest of the Gulf; it is influenced by inflow through the Strait of Belle Isle and is therefore not indicative of the climate in the rest of the Gulf. The CIL volume as defined by $T < 1^\circ\text{C}$ decreased significantly (to -1.5 SD) compared to 2014 conditions (-0.4 SD) and reached a volume similar to 2010 (-1.3 SD). The volume delimited by 0°C also decreased to a below-normal value (-0.9 SD).

The time series of the CIL regional average minimum core temperatures are shown in Figure 47. All regions, again except for Cabot Strait, show an increase in core temperature. The 2015 average temperature minimum (excluding the Magdalen Trough, the Strait of Belle Isle and the Magdalen Shallows) was 0.1°C, an increase of 0.2°C over 2014, and is shown in Figure 46 (bottom panel, green line). The overall 2015 CIL water mass properties are similar to observations of 2010.

NOVEMBER CIL CONDITIONS IN THE ST. LAWRENCE ESTUARY

The AZMP November survey provides a high-resolution conductivity-temperature-depth (CTD) sampling grid in the St. Lawrence estuary since 2006 although measurements are sparser in some years. This allows a higher resolution display of the CIL minimum temperature in the Estuary (Figure 48). The data also show the temporal warming (Figure 45) and thinning (Figure 47) of the CIL since the August survey. The CIL was warmer in November 2015 than in 2014, so much so that none of it was below 1°C. Figure 47 shows that the fairly rapid increase of the CIL minimum temperature occurring between August and November is fairly constant inter-annually in spite of the differences in August temperature.

GILBERT AND PETTIGREW (1997) CIL INDEX

The Gilbert and Pettigrew (1997) CIL index is defined as the mean of the CIL minimum core temperatures observed between 1 May and 30 September of each year, adjusted to 15 July. It was updated using all available temperature profiles measured within the Gulf between May and September inclusively since 1947 (black line of the bottom panel of Figure 46). As expected, the CIL core temperature interpolated to 15 July is almost always colder than the estimate based on August and September data for which no temporal corrections were made. This is because the CIL is eroded over the summer and therefore its core warms over time.

This CIL index for summer 2015 was -0.28°C, 0.3 SD above normal. The 0.21°C increase from the summer 2014 CIL index is consistent with the decrease in CIL volume between August 2014 and 2015 discussed above and the decrease of 0.2°C in the areal average of the minimum temperature in August. The warm winter conditions from 2010 to 2012 led to CIL indices that were still far below the record high observed in the 1960s and 1980s. The earlier CIL temperature minimums will be re-examined to confirm that they were calculated using data with sufficient vertical resolution to correctly resolve the core minimum temperature. It is also becoming increasingly clear that the winter mixed layer is not the only factor explaining

summertime CIL conditions and that mechanisms having a multi-year cumulative effect are required to explain the interannual autocorrelations observed. For example, this may be linked to temperatures below the CIL, which in warm years may create a higher temperature gradient that leads to higher heat fluxes and faster summertime CIL warming rates.

As a summary, Figure 49 shows selected time series of winter and summertime CIL conditions (June and September bottom temperatures also related to the CIL are outlined below) and highlights the strong correlations between these various time series. Surprisingly, the CIL had warmer-than-normal conditions in August whereas winter air temperatures were colder and the seasonal maximum sea-ice cover larger than normal. The August CIL was warmer and thinner in 2015 than in 2014 with conditions similar to those of 2010. The Gilbert & Pettigrew index which considers data from earlier in the season was near-normal and similar to the 2004 index.

MAGDALEN SHALLOWS JUNE SURVEY

A long-standing assessment survey covering the Magdalen Shallows has taken place in June for mackerel assessments and was since merged with the June AZMP survey. This survey provides good coverage of the temperature conditions that are greatly influenced by the cold intermediate layer that reaches the bottom at roughly half of the surface area at this time of time.

Near-surface waters warm quickly in June, mid-way between the winter minimum and the annual maximum in early August. This can introduce a bias if the survey dates are not the same each year. To account for this, the seasonal warming observed at the Shediac Valley AZMP monitoring station was evaluated. A linear regression was performed of temperature versus time for each meter of the water column for each year with monitoring data at Shediac Valley between May and July. Visual inspection showed that the depth-dependent warming rate was fairly constant for all years and an average was computed for every depth. Warming is maximal at the surface at 18°C per 100 days and, in spite of some uncertainties between 30 and 55 m, decreases almost proportionally with depth to reach 2°C per 100 days at 40 m, followed by a further linear decrease to reach 1°C per 100 days at 82 m (Galbraith and Grégoire 2015).

All available temperature profiles taken in June from a given year are binned at 1 m depth intervals (or interpolated if the resolution is too coarse) and then adjusted according to the sampling date to offset them to June 15th according to the depth-dependent warming rate extracted from Shediac Valley monitoring data. An interpolation scheme is used to estimate temperature at each 1 m depth layer on a 2 km resolution grid. Figure 50 shows temperatures and anomalies at depths of 20, 30 and 50 m. Figure 51 shows averages over the grids at 0, 10, 20, 30, 50 and 75 m for all years when interpolation was possible, as well as SST June averages since 1985, for both western and eastern regions of the Magdalen Shallows (Figure 22). Temperatures varied about the mean from below-normal to above-normal at 20, 30 and 50 m (Figure 50), as were the layer area averages (Figure 51).

BOTTOM WATER TEMPERATURES ON THE MAGDALEN SHALLOWS

Bottom temperature is also estimated at each point of the grids constructed from the June survey by looking up the interpolated temperature at the depth level corresponding to a bathymetry grid provided by the Canadian Hydrographic Service with some corrections applied (Dutil et al. 2012). The method is fully described in Tamdrari et al. (2012). A climatology was constructed by averaging all available temperature grids between 1981 and 2010 and anomaly grids were computed for each year. The June bottom temperature climatology as well as the 2015 reconstructed temperature and anomaly fields are shown in Figure 52. The same method was applied using the available CTD data from August and September, thus including the

multispecies surveys for the northern Gulf in August and for the Magdalen Shallows in September. These results are also shown in Figure 52. While much of the deeper bottom water temperatures are climatologically still below 0°C in June, a remnant from the winter near-freezing mixed layer that reached the bottom, most of the area usually warms to below 1°C by August-September. Temperature anomalies in coastal shallow waters range from <-2.5°C to >+2.5°C, but anomalies tend to be of smaller magnitude in deeper waters.

Time series of the bottom area covered by water in various temperature intervals were estimated from the gridded data for the June surveys as well as for the September multispecies survey on the Magdalen Shallows (Figure 53). The time series of areas of the Magdalen Shallows covered by water colder than 0, 1, 2, and 3°C in June and September are also shown in Figure 49 as part of the CIL summary. Unlike the very cold conditions observed on the bottom in 2008, only a small area of the bottom of the Magdalen Shallows was covered by water with temperatures <-1°C in June 2015; these are however colder conditions than during the 2010-2013 period when no bottom waters were that cold. The area covered by water temperatures <1°C in September had reached a low not seen since 1982 in 2012 but rebounded to near-normal in 2014 and 2015. Areas with $T < 0^{\circ}\text{C}$, $< 1^{\circ}\text{C}$, $< 2^{\circ}\text{C}$ and $< 3^{\circ}\text{C}$ were all near-normal in September 2015 (Figure 49). The thermograph network (Figure 29) also showed bottom temperatures at Shediac Valley (86 m) to be near normal (using a different reference period) for most of June to October.

DEEP WATERS (>150 M)

The deeper water layer (>150 m) below the CIL originates at the entrance of the Laurentian Channel at the continental shelf and circulates towards the heads of the Laurentian, Anticosti, and Esquiman channels without much exchange with the upper layers. The layer from 150 to 540 m is characterized by temperatures between 1 and >7°C and salinities between 32.5 and 35 (except for Mécatina Trough where near-freezing waters may fill the basin to 235 m in winter and usually persist throughout the summer). Interdecadal changes in temperature, salinity, and dissolved oxygen of the deep waters entering the Gulf at the continental shelf are related to the varying proportion of the source cold-fresh and high dissolved oxygen Labrador Current water and warm-salty and low dissolved oxygen slope water (McLellan 1957, Lauzier and Trites 1958, Gilbert et al. 2005). These waters travel from the mouth of the Laurentian Channel to the Estuary in roughly three to four years (Gilbert 2004), decreasing in dissolved oxygen from in situ respiration and oxidation of organic material as they progress to the channel heads. The lowest levels of dissolved oxygen (around 20 percent saturation in recent years) are therefore found in the deep waters at the head of the Laurentian Channel in the Estuary.

BOTTOM WATER TEMPERATURES IN AUGUST AND SEPTEMBER

The same method used to calculate bottom water temperature on the Magdalen Shallows was applied to the entire Gulf by combining all available CTD data from August and September, thus including the multispecies surveys for the northern Gulf in August and for the Magdalen Shallows in September into a single map (Figure 54). Most of the northeastern Gulf had above normal bottom temperatures, with large areas of Anticosti and Esquiman Channels above 6°C.

Time series of the bottom area covered by water in various temperature intervals were also estimated for the other regions of the Gulf based on August-September temperature profile data (Figures 55 and 56). Many areas had bottom waters colder than 0°C, and Mécatina Trough had bottom waters colder than -1°C. The figures also show compression of the bottom habitat area in the temperature range of 5-6°C in 1992. In 2012, a return of >6°C temperatures to the sea floor was observed. The bottom area covered by waters warmer than 6°C increased again in

2015 in Anticosti Channel, Esquiman Channel and Central Gulf, and reached again a record value in Anticosti Channel and Esquiman Channel while reducing bottom habitat area in the temperature range of 5–6°C.

The warm waters found at the bottom of Esquiman Channel and elsewhere are associated with the deep temperature maximum evident in the temperature profiles in these areas (e.g. Figure 3). The progression from a cold anomaly within the Gulf in 2009 to current conditions of the deep temperature maximum is shown on Figure 57. In 2012 and again in 2015, temperatures above 7°C were recorded in the Gulf near Cabot Strait. The Gulf-wide average and regional areal averages are shown in Figure 58. The deep maximum temperature Gulf-wide average was at a series record high in 2015, at 6.19°C, as were the regional averages for the northwest Gulf, Anticosti Channel and Esquiman Channel.

TEMPERATURE AND SALINITY MONTHLY MEANS

Monthly temperature and salinity averages were constructed for various depths using a method used by Petrie et al. (1996) but using the geographical regions shown in Figure 2. In this method, all available data obtained during the same month within a region and close to each depth bin are first averaged together for each year. Monthly averages from all available years and their standard deviations are then computed. This two-fold averaging process reduces the bias that occurs when the numbers of profiles in any given year are different. These monthly averages were further averaged into regional yearly time series that are presented in Figure 58 for 200 and 300 m. The 300 m observations in particular suggest that temperature anomalies are advected up-channel from Cabot Strait to the northwestern Gulf in two to three years, consistent with the findings of Gilbert (2004). The regional averages are weighted into a Gulf-wide average in accordance to the surface area of each region at the specified depth. These Gulf-wide averages are shown for 200 and 300 m in Figure 58 as well as for 150, 200, and 300 m in Figure 59. Linear trends in temperature and salinity at 300 m of 2.2°C and 0.3 per century, respectively are shown on Figure 59 (See also Galbraith et al. 2013 for other long term trends).

In 2015, the gulf-wide average salinities increased to above normal at all depths shown in Figures 58 and 59. Temperature at 200 m increased overall to reach a record-high of 5.6°C (+2.7 SD). It increased to regional record highs in the Northwest Gulf (5.2°C, +2.4 SD), Anticosti Channel (5.6°C, +2.8 SD), Esquiman Channel (5.9°C, +2.3 SD), and remained well above-normal in central Gulf (5.8°C, +2.5 SD, second highest) and Cabot Strait (6.2°C, +2.2 SD, third highest). Temperature at 250 m increased overall to reach 6.1°C (+2.8 SD); the highest on record. Temperature at 300 m increased overall to reach the record-high of 6.0°C (+3.4 SD). It increased to regional record highs in the Estuary (5.4°C, +1.8 SD), Northwest Gulf (5.8°C, +3.0 SD), Anticosti Channel (6.4°C, +3.2 SD), Esquiman Channel (6.3°C, +2.8 SD), and remained well above-normal in central Gulf (6.3°C, +3.2 SD, second highest) and Cabot Strait (6.2°C, +2.1 SD, third highest).

The warm anomaly present since 2010 at Cabot Strait (record in 2012) has been progressing up the channel towards the Estuary since then, but waters that have followed into the gulf have also remained very warm such that the average overall temperature may continue to increase (Figure 57).

DISSOLVED OXYGEN AND HYPOXIA IN THE ST. LAWRENCE ESTUARY

Figure 59 shows an update of the Gilbert et al. (2005) oxygen time series of the mean dissolved oxygen value at depths ≥ 295 m in the St. Lawrence Estuary. The value of $100 \mu\text{mol l}^{-1}$ corresponds approximately to 30% saturation, below which is considered to be hypoxic. Since some of the variability is associated with changing water masses, the temperature at 300 m in

the Estuary is also shown. The deep waters of the Estuary were briefly hypoxic in the early 1960s and have consistently been hypoxic since 1984. Dissolved oxygen decreased to its lowest annual average in 2015, at $54.5 \mu\text{mol l}^{-1}$ (-1.6 SD), corresponding to 18% saturation.

Based on interdecadal variability, the inflow of warmer waters to the Estuary is expected to deteriorate the hypoxic conditions since these waters are typically poorer in dissolved oxygen (McLellan 1957, Lauzier and Trites 1958, Gilbert et al. 2005). A change of dissolved oxygen of $147.4 \mu\text{mol l}^{-1}$ is accounted for by a 10.09°C temperature difference in source water masses in Gilbert et al. 2005, implying that a decrease of $1.46 \mu\text{mol l}^{-1}$ would be expected for each 0.1°C temperature increase (at Cabot Strait). Recent interannual variability in the Gilbert et al. 2005 updated time series did not show changes in dissolved oxygen that would have been expected to be associated with the increases in temperature. However during the last two years, the dissolved oxygen concentration decrease ($-11.7 \mu\text{mol l}^{-1}$) has been greater than expected considering the observed warming of Estuary bottom waters. The correlation between the temperature and oxygen timeseries shown on the bottom panel of Figure 63 explains 74% of the variance ($R^2 = 0.74$) and other factors can cause oxygen variability, such as interannual changes in the vertical flux of organic matter to the bottom waters of the Lower St. Lawrence Estuary.

SEASONAL AND REGIONAL AVERAGE TEMPERATURE STRUCTURE

In order to show the seasonal progression of the vertical temperature structure, regional averages are shown in Figures 60 to 63 based on the profiles collected during the March helicopter survey, the June AZMP and mackerel surveys, the August multi-species survey (September survey for the Magdalen Shallows), and the October-November AZMP survey. All additional archived CTD data for those months were also used, including a NOVA autonomous profiler deployed in June that was active until January 2016. The temperature scale was adjusted to highlight the CIL and deep-water features; the display of surface temperature variability is best suited to other tools such as remote sensing and thermographs. Average discrete depth layer conditions are summarized for the months of the 2014 and 2015 AZMP surveys in Figure 64 for temperature and in Figure 65 for salinity and 0-50 m stratification. For each survey the anomalies were computed relative to monthly temperature and salinity 1981-2010 climatologies calculated for each region, shown in grey as the mean value $\pm 0.5 \text{ SD}$ in Figures 60 to 63.

The March water temperature conditions are shown in Figure 60 but caution is needed in interpreting the profiles. Indeed, regional averaging of winter profiles does not work very well in the northeast Gulf (regions 3 and 4) because very different water masses are present in the area such as the cold Labrador Shelf intrusion with saltier and warmer deeper waters of Anticosti Channel or Esquiman Channel. For example, the sudden temperature increase near the bottom of Mécatina Trough in 2015 resulted from the deepest cast used in each of the averages, which contained warmer waters. Large changes near 200 m are due to our usual sampling cutoff near 200 m for the March airborne survey, with some casts being slightly deeper than others. The highlights of March water temperatures shown in Figure 60 include the previously discussed winter mixed layer, with below-normal temperatures and above-normal thickness. The thermocline was much shallower than usual in most regions, associated with temperatures well above normal at 200 m. Waters deeper than 100 m in Mécatina Trough were much warmer than normal, indicating no deep renewal from a Labrador Shelf intrusion.

Temperatures in June and August 2015 (Figures. 58 and 59 and summary in Figure 64) were characterized by CIL conditions that were below normal in thickness. The CIL temperature minimum was usually high in the Estuary. Deep-water temperatures were above normal in all

regions along the Laurentian Channel, with most regions showing increases compared with 2014 conditions in waters shallower than 250 m depth. Temperatures at the depth of the temperature maximum (200 to >250 m) remained above normal in Esquiman Channel and central Gulf, exceeding 6°C at depth, presumably advected in from the Cabot Strait recent record-high conditions. Deep waters in Mécatina Trough were colder in June than in March, indicating a later-than-usual intrusion from the Labrador Shelf, but returned to warmer conditions by August, then cold again in the fall (see also Figure 64). These changes indicate much more variability than is usually observed, with the warmer and saltier waters corresponding more to Esquiman Channel waters than typical Mécatina Trough waters. Waters beneath the CIL were much warmer and saltier than normal in the Estuary in November (Figure 63), indicating enhanced deep circulation into the Estuary.

CURRENTS AND TRANSPORTS

Currents and transports are derived from a numerical model of the Gulf of St. Lawrence, Scotian Shelf, and Gulf of Maine. The model is prognostic, i.e., it allows for evolving temperature and salinity fields. It has a spatial resolution of 1/12° with 46 depth-levels in the vertical. The atmospheric forcing is taken from the Global Environmental Multiscale (GEM) model running at the Canadian Meteorological Center (CMC). Freshwater runoff is obtained from observed data and the hydrological model, as discussed in the freshwater runoff section. A simulation was run for 2006–2015 from which transports were calculated. The reader is reminded that the results outlined below are not measurements but simulations and improvements in the model may lead to changes in the transport values.

Figures 66–68 show seasonal depth-averaged currents for 0–20 m, 20–100 m, and 100 m to the bottom for 2015. Currents are strongest in the surface mixed layer, generally 0–20 m, except in winter months when the 20–100 m and the 100 m to bottom averages are almost as high (note the different scale for this depth). Currents are also strongest along the slopes of the deep channels. The Anticosti Gyre is always evident but strongest during winter months, when it even extends strongly into the bottom-average currents.

Monthly averaged transports across seven sections of the Gulf of St. Lawrence are shown in Figure 69 for sections with estuarine circulation, and in Figure 70 for sections where only net transports are relevant. In Figure 69, the net transport integrates both up and downstream circulation and, for example, corresponds to freshwater runoff at the Pointe-des-Monts section. The outflow transport integrates all currents heading toward the ocean, while the estuarine ratio corresponds to the outflow divided by the net transports.

Transports through sections under the direct estuarine influence of the St. Lawrence River (e.g., Pointe-des-Monts) have a more direct response to change in freshwater runoff while others (e.g., Cabot Strait, Bradelles Bank) have a different response, presumably due to redistribution of circulation in the GSL under varying runoff. The estuarine circulation ratio is determined by the mixing intensities within the estuary and is greatly influenced by stratification. It is on average greatest during winter months and weakest during the spring freshet. In fact, it is sufficiently reduced in spring that the overall outward transport at Pointe-des-Monts reaches its minimum value in June even though this month corresponds to the third highest net transport of the year, i.e. the estuary becomes sufficiently stratified that fresh water runoff tends to slip on top of the denser salty waters underneath. In 2015, the estuarine ratio at Pointe-des-Monts was above-normal in late winter and early spring. The below-normal spring freshet led to above-normal Estuarine ratio and, perhaps paradoxically, above-normal outward transport at the Pointe-des-Monts section (e.g. enhanced estuarine entrainment).

HIGH FREQUENCY SAMPLING AZMP STATIONS

Sampling by the Maurice Lamontagne Institute began in 1991 at a station offshore of Rimouski (46° 40' N 68° 35' W, 320 m depth; Plourde et al. 2009), typically once a week during summer and less often during spring and fall and almost never in winter (Figure 71). In 2013, following several analyses that identified good correlations and correspondences between the prior AZMP Anticosti Gyre and Gaspé Current stations with the Rimouski station, it was decided to drop sampling efforts at these logistically difficult stations and integrate the Rimouski station officially in the AZMP program. The AZMP station in the Shediac Valley (47° 46.8' N, 64° 01.6' W, 84 m depth) is sampled on a regular basis by the Bedford Institute of Oceanography as well as occasionally by DFO Gulf Region and by the Maurice Lamontagne Institute during their Gulf-wide surveys (Figure 71). This station has been sampled irregularly since 1947, nearly every year since 1957, and more regularly during the summer months since 1999 when the AZMP program began. However, observations were mostly limited to temperature and salinity prior to 1999.

Isotherms and isohalines as well as monthly averages of layer temperature and salinity, stratification, and CIL core temperature and thickness at $<1^{\circ}\text{C}$ are shown for 2011-2015 for the Rimouski station in Figure 72 and for the Shediac Valley station in Figure 73. The oceanic climatologies are calculated from 1991-2010 data for Rimouski station, and for 1981-2015 for Shediac Valley (given the sparseness of data prior to 1999).

At the Rimouski station in 2015, the CIL typically had much below-normal thickness and above-normal minimum temperature, with higher anomalies than those prevailing in the Gulf. At 200-300 m, the gradual shift of cold-fresh waters present in 2010 to warmer-saltier waters advected from Cabot Strait lead to a shift to warm anomalies by May 2013 and series records in temperature (5.55°C) observed in September of 2015. The thermograph network also shows a Rimouski station series record of the daily average temperature (since 2006) of 5.57°C occurring in September, and a record-high monthly average of 5.52°C (Figures 27 and 33).

A mooring measuring temperature and salinity at multiple depths was deployed at Shediac Valley station in June and rotated in October such that the data from June to September are currently available. Because of the sparse sampling at this station, the monthly averages at 10, 20, 30 and 75 m of Figure 73 were calculated using the mooring data rather than the monitoring sorties data. This eliminated strong anomalies that would have been estimated for the shallower depths but did not alter the anomalies significantly at 75 m. Temperatures were generally near-normal at Shediac Valley station over the season.

Figure 74 shows the interannual variability of some bulk layer averages from May to October for the two stations. Bulk surface layer temperature, salinity and CIL minimum temperature were above normal at Rimouski station while stratification was below normal. Near-bottom temperature (290 m) was at a record high. Bulk surface layer (0-50 m) temperature and salinity, as well as bottom temperature, were above-normal at Shediac Valley station in 2015. Since these bulk layer averages are calculated from profile data, the mooring data were not used here and the results at Shediac Valley are likely biased. An automatic CTD is planned for deployment in summer 2016 and will yield much more reliable results.

OUTLOOK FOR 2015

Air temperatures were respectively 2.5°C and 2.1°C above-normal over the Gulf in January and February 2016, but below-normal by 1.4°C in March. This was the setting for the March 2016 survey, which provides an outlook for CIL conditions expected for the remainder of 2016. Figure 75 shows the surface mixed layer temperature, salinity, and thickness (at $T < 1^{\circ}\text{C}$ and $T < 0^{\circ}\text{C}$).

as well as the thickness and extent of the cold and saline layer that has intruded into the Gulf from the Labrador shelf. A large portion of the winter mixed layer was warm, above -1°C . The overall thickness and volume of the layer colder than -1°C was 5.6 SD below normal and the Cold Intermediate Layer for summer 2015 is therefore forecasted to be warmer than in 2013, with a Gilbert and Pettigrew (1997) index of around $+0.07^{\circ}\text{C}$ compared to -0.28°C in 2013.

Concerning deep waters, recall that record high temperatures were recorded in Cabot Strait in 2012 and again in 2014, and that overall the Gulf waters below 150 m were in 2015 at a 100-year record high. There were 14 stations sampled in March 2015 with some deep-water temperatures greater than 6°C , located from Cabot Strait up to Centre Gulf. One station had its deep temperature maximum reaching 7.17°C . This signifies the continuation of warmer than normal conditions at depth.

SUMMARY

Figure 75 summarizes SST, summertime CIL, and deep-water average temperatures. For May-November SST, a proxy is used prior to 1985 using the April-November air temperature anomaly averaged over all stations of Figure 4 except the two from the Estuary, similar to the index developed in Galbraith et al. (2012). Figure 76 shows a decrease from 2014 values for May-November average SST, but increases for the average temperature at 200 m and 300 m, as well as for the CIL temperature minimum. The SST was near-normal, temperature at 200 m and 300 m were the highest on record (5.6°C , $+2.7$ SD and 6.1°C , $+3.4$ SD). The CIL temperature minimum was above-normal ($+0.5$ SD) in contrast to the colder than normal winter air temperatures (-1.5°C , -1.0 SD).

Another summary of the temperature state of the Gulf of St. Lawrence over a shorter time span (since 1971) allows the inclusion of more data sets, and three sets of four time series are chosen to represent surface, intermediate and deep conditions (Figure 77). Here, sea-ice is grouped as an intermediate feature since all are associated with winter formation. Figure 77 shows the sums of these three sets of anomalies representing the state of different parts of the system and is reproduced on Figure 78 with each time series contribution shown as stacked bars (Pettit et al. 2007). These composite indices measure the overall state of the climate system with positive values representing warm conditions and negative representing cold conditions. The plot also indicates the degree of correlation between the various measures of the environment. In 2015, the surface index decreased to the 5th largest in the series, the intermediate index was near-normal and the deep index increased to a record high value.

KEY FINDINGS

- The annual average runoff from the St. Lawrence River measured at Québec City and Rivière was below-normal and near-normal in 2015 ($11\,100\,\text{m}^3/\text{s}$, -1.0 SD and $17\,000\,\text{m}^3/\text{s}$, $+0.1$ SD respectively). The spring freshet was below-normal at -1.1 SD and -0.5 SD, but timing was normal.
- The estuarine ratio at Pointe-des-Monts was above-normal in late winter and early spring. The below-normal spring freshet led to above-normal Estuarine ratio and, perhaps paradoxically, above-normal outward transport at the Pointe-des-Monts section.
- April-November air temperatures were normal with an anomaly of 0.0°C . Air temperatures in August were the second warmest on record (since 1873, $+2.2^{\circ}\text{C}$, $+2.8$ SD) when averaged over all stations, including series record highs at four stations.

- Winter (November-March) air temperatures were below normal by -1.2°C (± 1.5 SD), the coldest since 1985.
- Annual mean air temperature anomaly in 2015 was less normal over the Gulf (-0.4°C , ± 0.5 SD).
- The winter surface mixed water layer ($< 1^{\circ}\text{C}$) volume of 15 SD was 3.5 SD above the 1985-2015 average. The Labrador Shelf water intrusion into the Davis Strait was at a record low in March 2015, representing only 2% (± 1.5 SD) of the cold water in the Gulf.
- Ice ice maximum volume was slightly above normal ($+0.5$ SD) at 85 km² thick of 1 ice formed during the month of February, coldest since 1985.
- The August and intermediate water (IW) showed warmer ($+0.5$ SD) and thicker ($+1.5$ SD) for volume colder than 1°C than normal conditions in spite of the cold winter. The Gilbert & Perigue maximum temperature index, which includes data over a larger domain, was however near normal ($+0.3$ SD).
- In 2015, the timing of summer onset was near normal while peak season melting was later than normal ($+1.7$ weeks, ± 1.4 SD). The Eastern Northwest Gulf Arctost Channel and the Davis Strait had longer than normal seasons with surface temperatures above 10°C from $+0.5$ to $+1.5$ SD, about two weeks longer, while the rest were near normal.
- May-November sea surface temperatures averaged over the Gulf were generally below normal until July and above normal from in August and September 2015, leading to a near normal May-November average ($+0.2^{\circ}\text{C}$, ± 0.5 SD). Record highs were nevertheless reached in September averaged over the Gulf (17.18°C , above normal by $+1.2^{\circ}\text{C}$ and 3.5 SD) and in the following regions: February ($+1.2^{\circ}\text{C}$, $+2.5$ SD), Northwest Gulf ($+1.2^{\circ}\text{C}$, $+1.5$ SD), Arctost Channel ($+2.1^{\circ}\text{C}$, $+2.1$ SD) and the Davis Strait ($+1.2^{\circ}\text{C}$, $+2.5$ SD).
- Winter maxima in the Davis Strait were exchanged more frequently than usual, beginning with a reversal from a Labrador Shelf intrusion that came in only after the March survey.
- Deep water temperatures have been increasing in the Gulf, with record observations from Cabot Strait where temperature had reached a record high since 1970 in 2012 at 300 m. Temperature averaged over the Gulf at depths from 100 to 300 m all show record highs in 2015 and stand above 5°C at 200 and 300 m for the first time since 1970.
- Bottoms were covered by waters warmer than 5°C increased in 2015 in Arctost Channel, Esquimaux Channel and Central Gulf, and reached a record value in Arctost Channel and Esquimaux Channel while reducing its bottom habitat area in the temperature range of 5 - 6°C .

ACKNOWLEDGEMENTS

We are grateful to the people responsible for CTD data acquisition during the surveys used in this report:

- Groundwater station monitoring: Roger Flynn, Felix St-Pierre, Michel Gauthier, Sylvain Chartrand, Pierre Joly, Lys St-Amand, Lys St-Amand, Marie-Lys Dubé, Michel Gauthier, Yves Gagnon, René Desjardins, François Villeneuve, Caroline Lafleur, Jean-François St-Pierre, Michael St-Amand, Peter Gauthier, Bernard Pothier, Isabelle St-Pierre, Lucie Desrosiers.
- March 2015 helicopter survey: Peter Gauthier, René Desjardins, helicopter pilot Michel Dubé, ground technician Barry Corbett.

1. **Task 1:** Write a function `isPrime(n)` that takes an integer `n` and returns `True` if `n` is a prime number and `False` otherwise.
2. **Task 2:** Write a function `fibonacci(n)` that returns the `n`-th Fibonacci number.
3. **Task 3:** Write a function `isPalindrome(s)` that checks if a string `s` is a palindrome (reads the same forward and backward).
4. **Task 4:** Write a function `countVowels(s)` that counts the number of vowels in a string `s`.
5. **Task 5:** Write a function `mergeSort(arr)` that sorts an array `arr` using the merge sort algorithm.
6. **Task 6:** Write a function `findMax(arr)` that finds the maximum value in an array `arr`.
7. **Task 7:** Write a function `isAnagram(s1, s2)` that checks if two strings `s1` and `s2` are anagrams.
8. **Task 8:** Write a function `flatten(nestedList)` that flattens a nested list `nestedList` into a single list.
9. **Task 9:** Write a function `rotateImage(img)` that rotates a 2D image matrix 90 degrees clockwise.
10. **Task 10:** Write a function `calculateArea(shape, dimensions)` that calculates the area of a given shape (rectangle, triangle, circle) based on its dimensions.

FIGURE 1




FIGURE 1. The distribution of the Pacific halibut in the 1950s and 1960s. The distribution area is shaded in black. The maps show the coastline of the Pacific Northwest, with the distribution area located in the central and northern parts of the coast, extending from approximately 45°N to 55°N latitude and 125°W to 135°W longitude. The maps include latitude and longitude markings and a scale bar.



Figure 2. Gulf of St. Lawrence divided into macrographic regions. The grid cells are typically used in this model.




Figure 3. Typical seasonal progression of the bathy profiles of temperature observed in the Gulf of St. Lawrence. Profiles are averages of observations in August, June and November 2007 in the northern Gulf. The dashed line at 0°C shows a single winter temperature profile (March 2008), with near freezing temperatures in the top 75 m. The cold intermediate layer (CIL) is defined as the part of the water column that is colder than 7°C, although some authors use a different temperature threshold. Figures from Cabreret *et al.* (2012).




Figure 4. Monthly air temperatures and anomalies for 2014 and 2015 of selected stations around the Gulf as well as the average for all stations. The blue area represents the 1981–2010 climatological monthly mean ± 0.5 SD. Months with 4 or more days of missing data are omitted. The bottom scorecards are colour-coded according to the monthly normalized anomalies based on the 1981–2010 climatologies for each month, but the numbers are the monthly anomalies in $^{\circ}\text{C}$. For anomalies greater than 2 SD from normal, the prior year with a greater anomaly is indicated. Seasonal, December–March, April–November and annual anomalies are included for the all-station average.




Figure 5 Annual, April–November December–March mean air temperature anomalies averaged for the selected stations around the Gulf from Figure 4. The bottom scorecards are colour-coded according to the normalized anomalies based on the 1981–2010 climatology. Trends plus and minus their 95% confidence intervals are shown. April–November air temperature anomalies tend to be highly-correlated with May–November sea-surface temperature anomalies (Galbraith et al. 2012; Galbraith and Larouche 2013) whereas winter air temperature anomalies correlate highly with sea-ice cover parameters and winter mixed-layer volumes (Galbraith et al. 2010; Galbraith et al. 2013).




Figure 6. Monthly mean freshwater flow of the St. Lawrence River at Québec City (lower curve) and its sum with rivers flowing into the St. Lawrence Estuary (RIVSUM II, upper curve). The 1981–2010 climatological mean ($\pm 0.5 \text{ SD}$) is shown (blue shading). The scorecards are colour-coded according to the monthly anomalies normalized for each month of the year, but the numbers are the actual monthly anomalies in $10^3 \text{ m}^3 \text{ s}^{-1}$.




Figure 7. River discharge locations for the regional sums of runoffs listed in Table 2. Red and blue dots indicate rivers that have climatological mean runoff greater than $200 \text{ m}^3 \text{ s}^{-1}$ and between 100 and $200 \text{ m}^3 \text{ s}^{-1}$, respectively.




Figure 8. Monthly anomalies of the St. Lawrence River runoff and sums of all other major rivers draining into separate Gulf regions for 2014 and 2015. The scorecards are colour-coded according to the monthly normalized anomalies based on the 1981–2010 climatologies for each month, but the numbers are the monthly average runoffs in m³ s⁻¹. Numbers on the right side are annual climatological means. Runoff regulation is simulated for three rivers that flow into the Estuary (Saguenay, Manicouagan, Outardes).




Figure 9. Annual mean freshwater flow of the St. Lawrence River at Québec City and of the sum of all rivers flowing into regions of the Estuary and Gulf. The 1981–2010 climatological mean is shown as horizontal lines and indicated on the right side of the scorecards. Numbers in scorecards are normalized anomalies.




Figure 10. Locations of the Maurice Lamontagne Institute thermograph network stations in 2015, including oceanographic buoys that transmit data in real time (squares). Deep and shallow instruments are denoted by open circles and dots, while seasonal and year-round deployments are denoted by gray and black symbols. Shédiac station from DFO Gulf Region is also shown.




Figure 11. Sea-surface temperature climatological seasonal cycle in the Gulf of St. Lawrence. AVHRR temperature weekly averages for 1985 to 2010 are shown from May to November (ice-free months) for the entire Gulf (thick black line) and the cooler Lower St. Lawrence Estuary (thin black line), defined as the area west of the Pointe-des-Monts section and east of approx 69°30'W. Thermosalinograph data averages for 2000 to 2010 are shown for the head of the Laurentian Channel (at 69°30'W, grey dashed line) and for the average over the Gulf waters along the main shipping route between the Pointe-des-Monts and Cabot Strait sections (gray dash-dotted line). Monthly air temperature averaged over eight stations in the Gulf of St. Lawrence are shown offset by 2 weeks into the future (thick grey line; winter months not shown). Figure from Galbraith et al. (2012).




Figure 12. Thermosalinograph data at 8 m depth along the Montréal to St. John's shipping route: composite mean annual cycle of the water temperature for the 2000–2015 period (left panel), composite annual cycle of the water temperature for 2015 (middle panel), and water temperature anomaly for 2015 relative to the 2000–2015 composite (right panel).




Figure 13. Thermosalinograph near-surface temperature monthly anomalies for various sections along the main shipping lane. The numbers on the right are the 2000–2015 climatological means and standard deviations. The numbers in the boxes are normalized anomalies. The map shows all TSG data sampled in 2015. Those drawn in colour are within the main shipping corridor and are used in this report. Monthly average anomalies of temperatures measured close to the indicated blue section lines are shown in the other scorecard panels.




Figure 14. Sea surface temperature monthly averages for 2015 as observed with EarthSat remote sensing. Grey areas have no data for the period due to ice cover or clouds.




Figure 15. Low surface temperature monthly anomalies for 2018 based on monthly streamflow anomalies estimated by the 1986-2018 period observed with different stream gauging. They are often used as an index of streamflow anomalies due to increased snowmelt runoff.




Figure 10 displays the 100% $\hat{\Delta}_{\text{H}}^{\text{H}}(t)$ bootstrap resampling average and its half-spread for each region of the study. The data are generated by the 100% $\hat{\Delta}_{\text{H}}^{\text{H}}(t)$ stratigraphic resampling, $\text{mean} = 1.0$ (in the horizontal axis) and the spread according to the stratigraphic resampling procedure is the 100% $\hat{\Delta}_{\text{H}}^{\text{H}}(t)$ stratigraphic bootstrap spread. The values on the stratigraphic resampling procedure decrease.


[illegible]

Figure 3. (a) *Sty. thymus* 100 mg/kg BW, (b) *Sty. thymus* 500 mg/kg BW, and (c) *Sty. thymus* 1000 mg/kg BW. (d) *Sty. thymus* 100 mg/kg BW + *Sty. thymus* 500 mg/kg BW. (e) *Sty. thymus* 100 mg/kg BW + *Sty. thymus* 1000 mg/kg BW. (f) *Sty. thymus* 500 mg/kg BW + *Sty. thymus* 1000 mg/kg BW.




Figure 10. *Intermittent* and *continuous* sampling methods. *Intermittent* sampling involves taking samples at specific times and the probability of a given time being sampled is the same. *Continuous* sampling involves taking a 24-hr (or 24-hr) continuous sample of water from a well located in the middle of the sampling frequency (i.e., 12 hr). *Intermittent* and *continuous* sampling are identical in the case of a steady-state and *continuous* sampling is superior in the case of a non-steady-state system with a significant diurnal variation. The *intermittent* sampling is also superior.



Figure 1. A schematic diagram of the experimental setup. The diagram shows a cross-section of a system with a central region of interest. The central region is labeled 'Sample' and is surrounded by a 'Support' structure. The support structure is further divided into 'Inlet' and 'Outlet' sections. The diagram also shows a 'Flow' direction indicated by an arrow. The entire system is enclosed in a 'Container'.




Figure 24. Extended SST April to November monthly anomalies averaged over the Estuary (region 1 of the Gulf) and the Estuary subregion of the Saguenay – St. Lawrence Marine Park (RWSSSL), the proposed St. Lawrence Estuary Marine Protected Area (MPA), and Manicouagan WPA. The scorecards are colour-coded according to the monthly normalized anomalies based on the 1985–2110 climatologies for each month, but the numbers are the monthly average temperatures in °C. The 1985–2110 mean and standard deviation are indicated for each month on the right side of the table.



Figure 22. Areas defined as the western and eastern Magdalen Shallows.




Figure 23. AVHRR SST May to November monthly anomalies averaged over the Magdalen Shallows (region 8 of the Gulf) and the eastern and western subregions of the Magdalen Shallows. The scorecards are colour-coded according to the monthly normalized anomalies based on the 1985–2010 climatologies for each month, but the numbers are the monthly average temperatures in °C. The 1985–2010 mean and standard deviation are indicated for each month on the right side of the table.




Figure 24. Yearly number of weeks with mean weekly surface temperature $>10^{\circ}\text{C}$. (Top) Years with the minimum (1986, top left) and maximum (2006, top right) number of weeks are shown along with the 1985–2010 climatological average (lower left) and the chart for 2015. (Bottom) Anomaly for 2015 relative to 1985–2010 climatology, expressed in numbers of weeks.



Figure 25. Yearly number of weeks with mean weekly surface temperature $>10^{\circ}\text{C}$, averaged for the entire Gulf and each region of the Gulf. The scorecards are colour-coded according to the normalized anomalies based on the 1985–2010 time series, but the numbers are the average number of weeks above 10°C for each year.




Figure 26. Weekly average SST (1985-2015) matrices for the Gulf of St. Lawrence. Temperature anomaly proxies for the first and last occurrence of the 12°C isotherm based on June-July and September average air temperature are also shown (axes on right). Updated from Galbraith and Larouche 2013.

[illegible]

41

Lower North Shore / Basse Côte Nord




Figure 28. Thermograph network data. Daily mean 2015 temperatures compared with the daily climatology (blue areas are daily averages ± 0.5 SD, orange areas are daily average maximums ± 0.5 SD and bright blue areas are daily average minimums ± 0.5 SD) for stations of the lower north shore.

Southern Gulf / Sud du Golfe




Figure 29. Thermograph network data. Daily mean 2015 temperatures compared with the daily climatology (daily averages $\pm$ 0.5 SD; blue area) for stations of the southern Gulf. Data from 2014 are included if they were not all shown in last year's report. Red lines in the Île Shag panel span the historical dates when spring temperature increased over 1.5°C, a temperature associated with increased lobster mobility.





Figure 30. Thermograph network data. Daily mean 2015 salinities compared with the daily climatology (daily averages ± 0.5 SD; blue area) computed from all available stations.






Figure 10. Monthly mean temperatures at all stations from May 2014 to the October 2015 light survey. Temperature values are shown in 2014 and 2015. The number of days that each station was monitored is indicated in the top right. The color coding is according to the temperature gradient relative to the stratification of each station for each month. Numbers are monthly average temperatures with greater number of significant light included when average is lower.




Figure 10. Time series of the monthly average temperature recorded for selected stations of the meteorological network. The values within the rectangles are the temperature average values of the corresponding month within the year month. Rectangles are marked average temperatures. The black and white squares are indicated for each month at the right end of the figure.




Figure 1. The bird in flight, viewed from above. The bird is positioned in the center of each frame, with its wings spread. The background is a light, textured surface. The images are arranged in three rows and three columns, with a vertical line separating the first two columns from the third. The bird's head is pointed towards the top-left corner of the grid. The images show slight variations in the bird's wing position and the background texture, suggesting a sequence of frames or different lighting conditions.




Figure 8. Evolution of the vortex structure. The top row shows the initial state with a vortex core and surrounding flow. The bottom row shows the vortex core elongating and developing a secondary vortex. The left column is labeled $t = 0.0$ and the right column is labeled $t = 0.5$.



Figure 1. Absorbance spectrum of the solution of the polymer in THF. The inset shows the UV-Vis spectrum of the polymer in THF. The inset shows the UV-Vis spectrum of the polymer in THF. The inset shows the UV-Vis spectrum of the polymer in THF.




Figure 2. Molecular weight distribution of the polymer. The inset shows the GPC chromatogram of the polymer. The inset shows the GPC chromatogram of the polymer. The inset shows the GPC chromatogram of the polymer.




Figure 102: First and last day of ice occurrence, ice duration and maximum seasonal ice volume by region. The first when ice was first and last seen in days from the beginning of each year is indicated for each region, and the volume scale represents the annually based on the 1980–2012 climatology, with blue representing earlier first occurrence and later last occurrence. The threshold is 5% of the largest ice volume ever recorded in the region. Duration is the date for the date for the actual day of the year in volume, but the volume scaling is according to normalized anomalies based on the climatology of each region. Duration is the number of days that the threshold was exceeded. All results based on weekly data.




Figure 39 Seasonal maximum ice volume and area including the portion on the Scotian Shelf (excluding ice less than 15 cm thick), ice season duration and December-to-March air temperature anomaly (Figure adapted from Hammill and Galbraith 2012, but here not excluding small floes and adding February and March data to the air temperature anomalies). All sea-ice products are based on weekly data. Linear relations indicate losses of 17 km³, 30,000 km² and 14 days of sea-ice season for each 1°C increase in winter air temperature (R^2 of 0.74, 0.80 and 0.76 respectively).




Figure 40. Winter surface layer characteristics from the March 2015 helicopter survey compared with climatological means: surface water temperature (upper panel), temperature difference between surface water temperature and the freezing point (middle panel), and salinity (lower panel). Symbols are coloured according to the value observed at the station, using the same colour palette as the interpolated image. A good match is seen between the interpolation and the station observations where the station colours blend into the background. Black symbols indicate missing or bad data. The climatologies are based on 1996-2015 for salinity but excludes 2010 as an outlier for temperature and temperature above freezing.




Figure 41. Winter surface layer characteristics from the March 2015 helicopter survey compared with climatological means: estimates of the thickness of the Labrador Shelf water intrusion (upper panels), cold layer ($T < -1^{\circ}\text{C}$, $T < 0^{\circ}\text{C}$) thickness (middle panels), and maps indicating where the cold layer ($T < 0^{\circ}\text{C}$) reaches the bottom (in brown; lower panels). Station symbols are coloured according to the observed values as in Figure 40. For the lower panels, the stations where the cold layer reached the bottom are indicated with filled circles while open circles represent stations where the layer did not reach the bottom. Integrated volumes are indicated for the first six panels (including an approximation for the Estuary but excluding the Strait of Belle Isle). The climatologies are based on 1997-2015 for the Labrador Shelf water intrusion, 1996-2015 for the cold layer ($T < 0^{\circ}\text{C}$) but excludes 2010 for $T < -1^{\circ}\text{C}$.




Figure 42. Estimated volume of cold and saline Labrador Shelf water that flowed into the Gulf over the winter through the Strait of Belle Isle. The bottom panel shows the volume as a percentage of total cold-water volume ($< -1^\circ\text{C}$). The numbers in the boxes are actual values colour-coded according to their anomaly.




Figure 43. Left panel: winter surface cold ($T < -1^\circ\text{C}$ and $T < 0^\circ\text{C}$) layer volume (excluding the Estuary and the Strait of Belle Isle) time series (black and grey lines) and summer CIL index (blue dashed line). Right panel: Relation between summer CIL index and winter cold-water volume with $T < -1^\circ\text{C}$ (regression for 1996-2014 data pairs, excluding 1998 [see Galbraith 2006] as well as the 2010 and 2011 mild winters). Note that the CIL scale in the left panel is reversed.




Figure 44. Cold intermediate layer thickness ($T < 0^{\circ}\text{C}$, top panels; $T < 1^{\circ}\text{C}$, middle panels) and minimum temperature (bottom panels) in August and September 2015 (right) and 1985-2010 climatology (left). Station symbols are colour-coded according to their CIL thickness and minimum temperature. Numbers in the upper and middle panels are integrated CIL volumes and in the lower panels are monthly average temperatures.




Figure 45. Volume of the CIL colder than 0°C (blue) and colder than 1°C (red) in August and September (primarily region 8 in September). The volume of the CIL colder than 1°C in November for available years since 2006 is also shown for the St. Lawrence Estuary (dashed line).




Figure 48. CIL volume (top panel) delimited by IFC (in blue) and IFC (in red), and minimum temperature index (bottom panel) in the Gulf of St. Lawrence. The volumes are integrals of each of the annual interpolated thickness grids such as those shown in the top panels of Figure 44 excluding Mingana Trough and the Strait of Belle Isle. Rectangles on right side show mean $\pm$ 0.5 SD. In the lower panel, the black line is the updated Gilbert and Pettigrew (1997) index interpolated to 15 July (with dashed lines showing mean $\pm$ 0.5 SD) and the green line is the spatial average of each of the annual interpolated grid such as those shown in the two bottom panels of Figure 44, excluding Mingana Trough, the Strait of Belle Isle and the Magdalen Shallows. The numbers in the boxes are normalized anomalies relative to 1960-2010 climatologies constructed using all available years.




Figure 47. Temperature minimum of the CI, spatially averaged for the seven areas where the CI minimum temperature can be clearly identified. The spatial average of the November CI temperature minimum for available years since 2000 is also shown for the St. Lawrence Estuary (dashed line).




Figure 10. Sea surface temperature and salinity anomalies (°C and ‰) in March 1992. The 10° latitude/longitude grid is shown.




Figure 10. Distribution of various species. The top row shows the distribution of 'A. niger' in the area of the study. The middle row shows the distribution of 'A. niger' in the area of the study. The bottom row shows the distribution of 'A. niger' in the area of the study. The maps show the distribution of various species across a geographic area, with the top row showing the distribution of 'A. niger', the middle row showing 'A. niger', and the bottom row showing 'A. niger'.





Figure 16. A sequence of images showing a person's head and shoulders. The images are arranged in a 2x2 grid, with the top-left image being the most prominent. The images show a person's head and shoulders, with the person looking forward in the top row and looking down in the bottom row. The images are arranged in a 2x2 grid, with the top-left image being the most prominent.

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480	481	482	483	484	485	486	487	488	489	490	491	492	493	494	495	496	497	498	499	500	501	502	503	504	505	506	507	508	509	510	511	512	513	514	515	516	517	518	519	520	521	522	523	524	525	526	527	528	529	530	531	532	533	534	535	536	537	538	539	540	541	542	543	544	545	546	547	548	549	550	551	552	553	554	555	556	557	558	559	560	561	562	563	564	565	566	567	568	569	570	571	572	573	574	575	576	577	578	579	580	581	582	583	584	585	586	587	588	589	590	591	592	593	594	595	596	597	598	599	600	601	602	603	604	605	606	607	608	609	610	611	612	613	614	615	616	617	618	619	620	621	622	623	624	625	626	627	628	629	630	631	632	633	634	635	636	637	638	639	640	641	642	643	644	645	646	647	648	649	650	651	652	653	654	655	656	657	658	659	660	661	662	663	664	665	666	667	668	669	670	671	672	673	674	675	676	677	678	679	680	681	682	683	684	685	686	687	688	689	690	691	692	693	694	695	696	697	698	699	700	701	702	703	704	705	706	707	708	709	710	711	712	713	714	715	716	717	718	719	720	721	722	723	724	725	726	727	728	729	730	731	732	733	734	735	736	737	738	739	740	741	742	743	744	745	746	747	748	749	750	751	752	753	754	755	756	757	758	759	760	761	762	763	764	765	766	767	768	769	770	771	772	773	774	775	776	777	778	779	780	781	782	783	784	785	786	787	788	789	790	791	792	793	794	795	796	797	798	799	800	801	802	803	804	805	806	807	808	809	810	811	812	813	814	815	816	817	818	819	820	821	822	823	824	825	826	827	828	829	830	831	832	833	834	835	836	837	838	839	840	841	842	843	844	845	846	847	848	849	850	851	852	853	854	855	856	857	858	859	860	861	862	863	864	865	866	867	868	869	870	871	872	873	874	875	876	877	878	879	880	881	882	883	884	885	886	887	888	889	890	891	892	893	894	895	896	897	898	899	900	901	902	903	904	905	906	907	908	909	910	911	912	913	914	915	916	917	918	919	920	921	922	923	924	925	926	927	928	929	930	931	932	933	934	935	936	937	938	939	940	941	942	943	944	945	946	947	948	949	950	951	952	953	954	955	956	957	958	959	960	961	962	963	964	965	966	967	968	969	970	971	972	973	974	975	976	977	978	979	980	981	982	983	984	985	986	987	988	989	990	991	992	993	994	995	996	997	998	999	1000	1001	1002	1003	1004	1005	1006	1007	1008	1009	1010	1011	1012	1013	1014	1015	1016	1017	1018	1019	1020	1021	1022	1023	1024	1025	1026	1027	1028	1029	1030	1031	1032	1033	1034	1035	1036	1037	1038	1039	1040	1041	1042	1043	1044	1045	1046	1047	1048	1049	1050	1051	1052	1053	1054	1055	1056	1057	1058	1059	1060	1061	1062	1063	1064	1065	1066	1067	1068	1069	1070	1071	1072	1073	1074	1075	1076	1077	1078	1079	1080	1081	1082	1083	1084	1085	1086	1087	1088	1089	1090	1091	1092	1093	1094	1095	1096	1097	1098	1099	1100	1101	1102	1103	1104	1105	1106	1107	1108	1109	1110	1111	1112	1113	1114	1115	1116	1117	1118	1119	1120	1121	1122	1123	1124	1125	1126	1127	1128	1129	1130	1131	1132	1133	1134	1135	1136	1137	1138	1139	1140	1141	1142	1143	1144	1145	1146	1147	1148	1149	1150	1151	1152	1153	1154	1155	1156	1157	1158	1159	1160	1161	1162	1163	1164	1165	1166	1167	1168	1169	1170	1171	1172	1173	1174	1175	1176	1177	1178	1179	1180	1181	1182	1183	1184	1185	1186	1187	1188	1189	1190	1191	1192	1193	1194	1195	1196	1197	1198	1199	1200	1201	1202	1203	1204	1205	1206	1207	1208	1209	1210	1211	1212	1213	1214	1215	1216	1217	1218	1219	1220	1221	1222	1223	1224	1225	1226	1227	1228	1229	1230	1231	1232	1233	1234	1235	1236	1237	1238	1239	1240	1241	1242	1243	1244	1245	1246	1247	1248	1249	1250	1251	1252	1253	1254	1255	1256	1257	1258	1259	1260	1261	1262	1263	1264	1265	1266	1267	1268	1269	1270	1271	1272	1273	1274	1275	1276	1277	1278	1279	1280	1281	1282	1283	1284	1285	1286	1287	1288	1289	1290	1291	1292	1293	1294	1295	1296	1297	1298	1299	1300	1301	1302	1303	1304	1305	1306	1307	1308	1309	1310	1311	1312	1313	1314	1315	1316	1317	1318	1319	1320	1321	1322	1323	1324	1325	1326	1327	1328	1329	1330	1331	1332	1333	1334	1335	1336	1337	1338	1339	1340	1341	1342	1343	1344	1345	1346	1347	1348	1349	1350	1351	1352	1353	1354	1355	1356	1357	1358	1359	1360	1361	1362	1363	1364	1365	1366	1367	1368	1369	1370	1371	1372	1373	1374	1375	1376	1377	1378	1379	1380	1381	1382	1383	1384	1385	1386	1387	1388	1389	1390	1391	1392	1393	1394	1395	1396	1397	1398	1399	1400	1401	1402	1403	1404	1405	1406	1407	1408	1409	1410	1411	1412	1413	1414	1415	1416	1417	1418	1419	1420	1421	1422	1423	1424	1425	1426	1427	1428	1429	1430	1431	1432	1433	1434	1435	1436	1437	1438	1439	1440	1441	1442	1443	1444	1445	1446	1447	1448	1449	1450	1451	1452	1453	1454	1455	1456	1457	1458	1459	1460	1461	1462	1463	1464	1465	1466	1467	1468	1469	1470	1471	1472	1473	1474	1475	1476	1477	1478	1479	1480	1481	1482	1483	1484	1485	1486	1487	1488	1489	1490	1491	1492	1493	1494	1495	1
---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	---






Figure S3. Time series of the bottom areas covered by different temperature bins in June (top) and August-September (bottom) for the Magdalen Shallows (region 8). Data are mostly from September for the bottom panel.




Figure 54. August-September bottom temperature climatology (top), 2015 observations (middle) and anomaly (bottom).




Figure 55. Time series of the bottom areas covered by different temperature bins in August and September for regions 1 to 4. The panels are separated by a black horizontal line into shallow (<100m) and deep (>100 m) areas to distinguish between warmer waters above and below the CIL. The shallow areas are shown on top using the area scale on the right-hand side and have warmer waters shown starting from the top end. The deep areas are shown below the horizontal line and have warmer waters shown starting at the bottom end. The CIL areas above and below 100 m meet near the horizontal line.




Figure 56. Time series of the bottom areas covered by different temperature bins in August and September for regions 5 to 7. The panels are separated into shallow (<100 m) and deep (>100 m) areas to distinguish between warmer waters above and below the CIL. See Figure 55 caption.




Figure 57. Map of the deep temperature maximum found typically between 200 and 300 m, 2008-2015. Maps are interpolated from August-September data available for each year.




Figure 58. Deep layer temperature, salinity, and dissolved oxygen. Gulf averages for temperature and salinity are shown for 150, 200, 250, 300 m, as well as for the deep temperature maximum usually found between 200 and 300 m. Regional averages are shown for 200 and 300m, and deep temperature maximum. Dissolved oxygen is shown with an inverted colour scheme for the updated Gilbert et al. 2005 time series as well as recent regional averages at 300 m obtained using a Seabird SBE43 CTD sensor. The numbers on the right are the 1981–2010 climatological means and standard deviations. The numbers in the boxes are normalized anomalies, except for oxygen where physical values are shown for easier comparison between the Gilbert et al. 2005 time series and the CTD data.




Figure 59. Layer-averaged temperature and salinity time series for the Gulf of St. Lawrence and dissolved oxygen between 295 m and the bottom in the deep central basin of the St. Lawrence Estuary. The temperature and salinity panels show the 150 m, 200 m, and 300 m annual averages and the horizontal lines are 1981–2010 means. Sloped lines show linear regressions for temperature and salinity at 300 m of respectively 2.2°C and 0.3 per century. The horizontal line in the oxygen panel corresponds roughly to 30% saturation and marks the threshold of hypoxic conditions. In addition to the dissolved oxygen time series (light blue), the lower panel also shows temperature (dark blue) at 300 m in the Estuary.

March/mars 2015




Figure 60. Mean temperature profiles observed in each region of the Gulf during the March helicopter survey. The shaded area represents the 1981–2010 (but mostly 1980–2010) climatological monthly mean $\pm$ 1 S.D. Mean profiles for 2014 are also shown for comparison.

June/July 2015




Figure 21. Mean temperature profiles observed in each region of the Gulf during June. The shaded area represents the 1981-2010 climatological monthly mean ± 2 SD. Mean profiles for 2014 are also shown for comparison.

August-September 2015




Figure 10. Mean temperature profiles observed at each region of the Gulf during August and September 2015. The dashed line represents the 1981-2010 climatological mean. (a-h) are the August to August 1 through 7 and the September to August 8. Mean profiles for 2015 are also shown for comparison.




Figure 10. Mean temperature profiles (depth) in upper 200m of the water column for October, November, and December. The profiles were generated by the 1991-2000 depth-averaged monthly mean water temperature profiles (1991-2000) and the 1991-2000 standard deviation.



Figure 10. Distribution of various bird species in the United States. The maps show the distribution of various bird species in the United States. The maps show the distribution of various bird species in the United States. The maps show the distribution of various bird species in the United States.



Figure 1. Scatter plots showing the relationship between various variables. The plots are arranged in a 4x2 grid. Each plot has a title and axes. The plots show different distributions of data points, with some showing a clear positive or negative correlation and others showing a more complex relationship.




Figure 68. Depth-averaged currents from 0 to 20 m for each three-month period of 2015.




Figure 67. Depth-averaged currents from 20 to 100 m for each three-month period of 2015.




Figure 68. Depth-averaged currents from 100 m to the bottom for each three-month period of 2015.




Figure 69. Monthly averaged modelled transports and estuarine ratio across sections of the Gulf of St. Lawrence since 2006. The numbers on the right are the 2006–2015 means and standard deviations. The numbers in the boxes are normalized anomalies. Colours indicate the magnitude of the anomaly. Sv (Sverdrup) are units of transport equal to $10^6 \text{ m}^3 \text{ s}^{-1}$.




Figure 70. Mixed-layer averaged mixed transports across sections of the Gulf of St. Lawrence since 2005. The numbers on the right are the 2006-2013 means and standard deviations, with positive values toward east and north. The numbers in the boxes are normalized anomalies. Colours indicate the magnitude of the anomaly (e.g., negative anomalies are still shown in red when the mean transport is negative across the section).




Figure 75. Sampling frequency and positions of the A2000 stations Remoudeu and Shadow Valley.




Figure 15. Infrared and Raman spectra of the DMSO-water and water-ice systems. The infrared spectra and Raman spectra were recorded according to the procedure described in the text. (a) pure DMSO, (b) 10% water, (c) 20% water, (d) 30% water, (e) pure water. The peaks of the ice-like network are labeled: (a) ice-like network, (b) ice-like network, (c) ice-like network, (d) ice-like network, (e) ice-like network.

Shedden Valley Temperature




Figure 10. Shedden Valley Temperature. The top row shows vertical temperature profiles for the years 1981 through 1985. The bottom row shows cross-sectional temperature maps for the years 1981 through 1985. The color scale on the right indicates temperature values from 0 to 100.




Figure 10. Heatmap of errors (corrected and corrected) for the 'Correct' and 'Wrong type' categories. The x-axis represents the error type, and the y-axis represents the error value. The 'Correct' heatmap shows a high concentration of 'No error' for the '1000000' value, while the 'Wrong type' heatmap shows a high concentration of 'Wrong type' for the '1000000' value.





Figure 1. A series of eight line drawings illustrating the sequence of head and neck movements during a single cycle of a bird's head and neck movement. The drawings are arranged in two columns, with the left column showing a sequence of four drawings and the right column showing a sequence of four drawings. The drawings are labeled with numbers 1 through 8, indicating the sequence of the movement.






Figure 77 Surface and daily induction rate for various compounds (Figure 75).




Figure 7B. Composite climate indices (white lines and dots) derived by summing various normalized anomalies from different parts of the environment (colored boxes stacked above the abscissa are positive anomalies, and below are negative). Top panel sums anomalies representing the entire water column, middle panel sums shallow anomalies and bottom.